

ON THERMODYNAMICS OF MULTISPECIES ANYONS

Serguei B. Isakov*,
 Stefan Mashkevich^{**,† 1} and Stéphane Ouvry^{† 2}

* Medical Radiology Research Center, Obninsk, Kaluga Region 249020, Russia

** Institute for Theoretical Physics, Kiev 252143, Ukraine

† Division de Physique Théorique³, IPN, Orsay Fr-91406

Abstract: We address the problem of multispecies anyons, i.e. particles of different species whose wave function is subject to anyonlike conditions. The cluster and virial coefficients are considered. Special attention is paid to the case of anyons in the lowest Landau level of a strong magnetic field, when it is possible (i) to prove microscopically the equation of state, in particular in terms of Aharonov-Bohm charge-flux composite systems, and (ii) to formulate the problem in terms of single-state statistical distributions.

IPNO/TH 95-21

March 1995

¹*email: mash@phys.unit.no*

²*and LPTPE, Tour 12, Université Paris 6 / email: OUVRY@IPNCLS.IN2P3.FR*

³*Unité de Recherche des Universités Paris 11 et Paris 6 associée au CNRS*

1 Introduction

Considering quantum mechanical indistinguishable particles, one uses to speak about their “statistical” (or “exchange”) interaction. It has no classical analogue, since there is no classical interaction force, but it influences the many-particle spectrum. Under usual circumstances, with bosons and fermions only, the rules of combining single-particle states for indistinguishable particles are affected, while particles of different species remain independent. For anyons [1, 2], however, this interaction is nontrivial, since it leads to the absence of single-particle states, and even more, to the possibility that particles of different species can become mutually “entangled”; hence the problem of multispecies anyons demands special consideration.

It is also believed that this problem may be of direct physical interest, particularly in connection with the $t - J$ model, which might be relevant to the high- T_c superconductivity. The question about statistics of excitations therein, spinons and holons, is still unsettled, and it is anticipated that their “mutual statistics” may be nontrivial [3]. Another possible application is the layered Hall media where FQHE quasiparticles in different layers can be viewed as anyons with mutual statistical interactions [4]. Finally, the same situation arises in the usual quantum Hall liquid: In a state where exactly m Landau levels are filled, the excitations are characterized by “charge vectors” of dimension m and there are different nontrivial phase factors associated with interchanging different sorts of excitations [5].

In this paper we address certain points concerning thermodynamics of multispecies anyons, by starting from microscopic considerations on the wave functions. For the sake of simplicity, we do the calculations for two species only, but a generalization to more than two species would be straightforward. The fugacity expansion of the thermodynamic potential now involves two fugacities, and the equation of state, respectively, two densities; hence, the cluster and virial coefficients are labeled by two indices. The mixed virial

coefficient of the lowest order, a_{11} , is trivially calculable, but exact expressions for higher-order ones are certainly not easier to obtain than in the one-species case. To proceed further, we investigate the case when all the particles are in the lowest Landau level of a strong magnetic field, and, using microscopic arguments as in [6], we obtain the equation of state which coincides with the one conjectured in [7] and derived in [8] from a definition of fractional statistics à la Haldane [9] (exclusion statistics), using a state-counting argument. A mean-field interpretation of this equation is also proposed. Also, in this (and only in this) case it becomes possible to interpret the system in terms of statistical mechanics of a gas of free particles, by following essentially the same lines as in [10]. A statistical distribution is derived, which again coincides with the one for exclusion statistics [8, 11]. We summarize the expressions for the cluster and virial coefficients of a multispecies system, in the Appendix.

2 Mutual fractional exchange statistics

When particles are distinguishable, their wave functions have no definite behavior with respect to their interchange, but one can impose certain conditions that they should satisfy under a double interchange, or, equivalently, a winding of one particle around the other one. In three dimensions, as usually, wave functions are single-valued, whereas in two dimensions a non-trivial phase factor may arise. In fact, if there are M species of particles, one has an $M \times M$ matrix α_{ab} of statistical parameters, such that the wave function of the whole system picks up a phase

- $\exp[i\pi\alpha_{aa}]$ under an *interchange* of two particles of species a
- $\exp[2i\pi\alpha_{ab}]$ under *winding* a particle of species a around a particle of species b

(provided, in both cases, that the complete closed path encloses no other particles). The second condition, of course, holds for $a = b$ as well, but the first one in this case is stronger. Note that there is periodicity in α_{aa} 's with period 2, but in α_{ab} 's with period 1. Note also that since “winding particle a around particle b ” and “winding particle b around particle a ” is one and the same operation, in the system of their center of mass, the matrix α_{ab} must be symmetric.

The above condition on the wave function can also be understood within the approach where statistics appear as kinematic properties of the configuration space [1]. The configuration space for a system containing M species of identical particles can be constructed in analogy with the case of a single species [12]. Irreducible representations of the fundamental group of the configuration space determine properties of the wave function under exchange of particles. In [12] it was found that the fundamental group for the configuration space for a system with M species of identical particles is $B_{N_1 \dots N_M}$, a generalization of the braid group B_N . One-dimensional irreducible representations of $B_{N_1 \dots N_M}$ (corresponding to spinless particles) are labeled by M parameters associated with exchange of particles of the same species (which may be identified with the above α_{aa}) and $\frac{1}{2}M(M-1)$ parameters associated with exchange of particles of different species (which may be identified with the above α_{ab}) [12].

One important observation is now in order. A physical picture usually associated with anyons is the model of charge-flux composites [2], where the origin of the anyonic interchange phase is the Aharonov-Bohm interaction of the charges and fluxes ascribed to the particles. In the multispecies case, each species a would be characterized by a charge e_a and a flux ϕ_a . Now, if one writes the Hamiltonian in the regular gauge, including the relevant vector potentials, it turns out that a singular gauge transformation to the free

Hamiltonian (anyon gauge) is possible only if the equality

$$e_a \phi_b = e_b \phi_a \quad (1)$$

holds, for each a and b . In this case, one will have

$$\alpha_{ab} = e_a \phi_b / 2\pi, \quad (2)$$

and the matrix α_{ab} is indeed symmetric. Equations (1) and (2) together imply that

$$\alpha_{ab} = \pm \sqrt{\alpha_{aa} \alpha_{bb}}. \quad (3)$$

Thus, only the diagonal statistical parameters are independent, save for the sign. However, a situation may be imagined when there are several Aharonov-Bohm gauge fields. Eq. (1) then becomes

$$e_a^\beta \phi_b^\beta = e_b^\beta \phi_a^\beta, \quad (4)$$

β labeling the gauge fields, while (2) becomes $\alpha_{ab} = \sum_\beta e_a^\beta \phi_b^\beta / 2\pi$, and (3) no longer holds. Therefore, one may consider all α_{ab} 's as independent, but in the Aharonov-Bohm model this would, generally speaking, demand to have several gauge fields. Note at the same time that in a model where the fluxes appear as a result of coupling the particles to Chern-Simons gauge fields [13], (4) is satisfied automatically.

Consider now the simplest case of $M = 2$, and denote for brevity $\alpha_{11} = \alpha_1$, $\alpha_{22} = \alpha_2$, $\alpha_{12} = \gamma$. A many-particle wave function of a system of N_1 particles of species 1 and N_2 particles of species 2 that satisfies the above-discussed interchange conditions reads in the anyon gauge

$$\Psi = \prod_{j < k}^{N_1} (z_j - z_k)^{\alpha_1} \prod_{m < n}^{N_2} (\zeta_m - \zeta_n)^{\alpha_2} \prod_{j, m} (z_j - \zeta_m)^\gamma \Phi(\{z_j\}, \{\zeta_m\}) \quad (5)$$

$\alpha_1, \alpha_2, \gamma > 0$ is meant, while the replacement $(z_j - z_k)^{\alpha_1} \rightarrow (z_j^* - z_k^*)^{-\alpha_1}$ is to be made if $\alpha_1 < 0$, etc. Φ is a single-valued function symmetric with respect to the coordinates

$\{z_j\}$ (species 1) as well as with respect to the coordinates $\{\zeta_m\}$ (species 2), but arbitrary with respect to $z \leftrightarrow \zeta$. This is the starting point for constructing the $(N_1 + N_2)$ -particle spectrum. Obviously, as far as γ is not an integer, all $(N_1 + N_2)$ coordinates are “entangled” together, while if γ is integer, the problem can be factorized into the one of N_1 particles and the one of N_2 particles.

Knowing the spectrum, one may find the partition function $Z_{N_1 N_2} = \text{Tr} \exp(-\beta H_{N_1 N_2})$, then proceed to the grand partition function

$$\Xi = \sum_{N_1, N_2} z_1^{N_1} z_2^{N_2} Z_{N_1 N_2}, \quad (6)$$

where $z_a = \exp(\beta \mu_a)$ are the fugacities, in order to obtain the cluster expansion

$$\ln \Xi = \sum_{k_1, k_2} b_{k_1 k_2} z_1^{k_1} z_2^{k_2} \quad (7)$$

and the virial expansion for the equation of state

$$\beta P = \sum_{k_1, k_2} a_{k_1 k_2} \rho_1^{k_1} \rho_2^{k_2} \quad (8)$$

(see Appendix).

Clearly, for the virial coefficients a_{n0} and a_{0n} one has

$$a_{n0} = a_n(\alpha_1), \quad a_{0n} = a_n(\alpha_2), \quad (9)$$

where $a_n(\alpha)$ is the n -th virial coefficient of anyons with statistical parameter α . Among these, only a_2 is known exactly [14],

$$a_2(\alpha) = \left[\frac{1}{4} - \frac{1}{2}(1 - \alpha)^2 \right] \lambda^2, \quad (10)$$

with $\lambda = \sqrt{2\pi\beta/m}$. The mixed second-order coefficient, a_{11} , involves the $N_1 = N_2 = 1$ case. Eq. (5) then turns into

$$\Psi = (z - \zeta)^\gamma \Phi(z, \zeta), \quad (11)$$

where Φ is *any* single-valued function. Since it may always be represented as a sum of symmetric and antisymmetric functions, the partition function Z_{11} can be written as

$$Z_{11} = Z_2(\gamma) + Z_2(1 + \gamma), \quad (12)$$

$Z_2(\gamma)$ being the 2-anyon partition function with statistical parameter γ . Using the formulas from the Appendix, we get

$$a_{11} = a_2(\gamma) + a_2(1 + \gamma) = \gamma(1 - \gamma)\lambda^2. \quad (13)$$

For $\gamma = 0, 1$, this coefficient vanishes, reflecting the fact that particles of different species are mutually independent.

3 Quantum and statistical mechanics in the lowest Landau level

Calculating higher-order virial coefficients would demand solving a multiparticle problem, which is in general impossible. In the one-species case, however, -when the system is projected in the lowest Landau level (LLL) of a strong external magnetic field, the whole virial expansion can be obtained[6]. Note that for the gap above the ground state to be bounded from below and thus the projection onto the LLL to be justified, the particle fluxes must be antiparallel to the external field, for example if $B < 0$, one should have $\phi > 0$. If the particle charge e is taken to be positive, then the necessary condition is $\alpha \in [0, 1]$ ⁴. The equation of state then is

$$\beta P = \rho_L \ln \left(1 + \frac{\nu}{1 - \alpha\nu} \right) \quad (14)$$

⁴Note that this convention for signs is opposite to [6, 7] and coincides with [8].

($\rho_L = m\omega_c/\pi$ is the Landau degeneracy per unit area, $\rho = N/V$ is the density, $\nu = \rho/\rho_L$ is the filling factor, $\omega_c = |eB|/2m$ is half the cyclotron frequency; “strong magnetic field” means $\beta\omega_c \gg 1$).

It has been shown [8] that the same equation (14) is obtained from Haldane’s definition of fractional statistics [9], usually referred to as exclusion statistics, with α being the quantity by which the number of available single-particle states diminishes as one particle is added to the system. Why the equations coincide can be understood within a mean-field approximation (where it becomes possible to speak about single-particle states): Since the anyon fluxes are antiparallel to the external field, the flux $\phi = 2\pi\alpha/e$ of a particle added partially screens the external flux $\Phi = BV$ and thus diminishes the number of states available for other particles; when N particles are added, the Landau level degeneracy per unit area becomes $\rho_L - \rho\alpha$.

A multispecies generalization of (14) within the framework of exclusion statistics is as follows: If α_{ab} is the decrease in the number of single-particle states available for a -th species caused by addition of one particle of b -th species, then

$$\beta P = \rho_L \sum_a \ln \left(1 + \frac{\nu_a}{1 - \sum_b \alpha_{ab}\nu_b} \right), \quad (15)$$

with $\nu_a = \rho_a/\rho_L$ [7, 8].

The physical picture would be the following: a particle of a -th species carries statistical charges e_a^β , which couple to the statistical gauge fields, and an electric charge e which couples to the external magnetic field and has to be common to all species. In fact, we are forced to restrict ourselves to the Landau density ρ_L being the same for all species (whereas in [7, 8], where mean field arguments are used, different Landau densities ρ_{La} are possible), because it is only in this case that exact eigenstates can be constructed.

In the two-species case, by using microscopic arguments for the LLL spectrum, we will

now show that the above equation indeed holds with α_{ab} 's being the mutual statistical parameters defined previously: it is then almost certain that (15) is correct for any number of species.

If there is only one statistical charge, then, as already emphasized, γ is not independent from the α_{aa} 's. Now, both ϕ_1 and ϕ_2 have to be positive, and it follows then from (1) that e_1 and e_2 must have the same sign; choosing it again to be positive, we have $\alpha_1, \alpha_2 \in [0, 1]$, and, by virtue of (2), γ is positive as well, so the constraint (3) would read

$$\gamma = \sqrt{\alpha_1 \alpha_2}. \quad (16)$$

However, our considerations will hold in the case of several gauge fields as well, and the result will be true for arbitrary $\gamma \in [0, 1]$ (with the condition, however, that the gap above the ground state be bounded from below by a quantity of the order the Landau gap $2\omega_c$, which is certainly true for γ small enough).

By analogy with [6], we add a harmonic attraction $\sum_{j=1}^{N_1+N_2} \frac{1}{2} m \omega^2 r_j^2$ and observe, from (5), that the spectrum of the $(N_1 + N_2)$ -particle system in the LLL becomes that of a system of N_1 and N_2 bosons (mutually *independent*) in a “harmonic-Landau” potential whose single-particle spectrum reads

$$\varepsilon_\ell = \omega_t + (\omega_t - \omega_c)\ell \quad (17)$$

($\omega_t = \sqrt{\omega_c^2 + \omega^2}$, $\ell = 0, 1, \dots$), plus a constant shift

$$\Delta E_{N_1 N_2}(\alpha_1, \alpha_2, \gamma) = \left[\frac{N_1(N_1 - 1)}{2} \alpha_1 + \frac{N_2(N_2 - 1)}{2} \alpha_2 + N_1 N_2 \gamma \right] (\omega_t - \omega_c) \quad (18)$$

coming from the anyonic prefactor in (5). One then has

$$Z_{N_1 N_2} = e^{-\beta \Delta E_{N_1 N_2}(\alpha_1, \alpha_2, \gamma)} Z_{N_1}^b Z_{N_2}^b, \quad (19)$$

where

$$Z_N^b = \frac{e^{-N\beta\omega_t}}{(1 - e^{-\beta(\omega_t - \omega_c)})(1 - e^{-2\beta(\omega_t - \omega_c)}) \dots (1 - e^{-N\beta(\omega_t - \omega_c)})} \quad (20)$$

is the N -boson partition function in the harmonic-Landau potential.

For the cluster coefficients, one gets (see Appendix for the prescription for passing to the thermodynamic limit)

$$b_{k_1 0} = \rho_L V \frac{1}{k_1} \prod_{l_1=1}^{k_1-1} \left(1 - \frac{k_1 \alpha_1}{l_1}\right) \cdot e^{-k_1 \beta \omega_c}, \quad (21)$$

$$b_{k_1 k_2} = -\rho_L V \gamma \frac{(k_1 + k_2)}{k_1 k_2} \prod_{l_1=1}^{k_1-1} \left(1 - \frac{k_1 \alpha_1 + k_2 \gamma}{l_1}\right) \prod_{l_2=1}^{k_2-1} \left(1 - \frac{k_2 \alpha_2 + k_1 \gamma}{l_2}\right) \cdot e^{-(k_1 + k_2) \beta \omega_c}, \quad (22)$$

and $b_{0k_2} = b_{k_2 0}|_{\alpha_1 \leftrightarrow \alpha_2}$. Note that, as expected, (22) does not reduce to (21) when $k_2 = 0$ (in particular, $b_{k_1 k_2}$ certainly vanishes for $\gamma = 0$, while $b_{k_1 0}$ is γ independent). The virial coefficients calculated herefrom take the form ($\beta \omega_c \gg 1$)

$$a_{k_1 k_2} = -\frac{1}{\rho_L^{k_1 + k_2 - 1}} \frac{(k_1 + k_2 - 1)!}{k_1! k_2!} \left\{ [(\alpha_1 - 1)^{k_1} - \alpha_1^{k_1}] \gamma^{k_2} + [(\alpha_2 - 1)^{k_2} - \alpha_2^{k_2}] \gamma^{k_1} \right\}, \quad (23)$$

and the equation of state then is

$$\beta P = \rho_L \left[\ln \left(1 + \frac{\nu_1}{1 - \alpha_1 \nu_1 - \gamma \nu_2} \right) + \ln \left(1 + \frac{\nu_2}{1 - \alpha_2 \nu_2 - \gamma \nu_1} \right) \right], \quad (24)$$

which indeed coincides with (15).

A mean-field interpretation is possible in this case as well. Indeed, what figures in the Hamiltonian for a -th species is the sum $e\vec{A} + \sum_{\beta} e_a^{\beta} \vec{A}^{\beta}$, where $\vec{A}$ is the vector potential of the magnetic field and $\vec{A}^{\beta}$ is the β -th gauge field potential created by all other particles. If *all* the gauge fields are averaged, so that $B^{\beta} = \sum_b N_b \phi_b^{\beta} / V = \sum_b \rho_b \phi_b^{\beta}$, then the Landau level degeneracy per unit area becomes $\frac{1}{2\pi} (|eB| - \sum_{\beta} e_a^{\beta} B^{\beta}) = \rho_L - \sum_b \rho_b \alpha_{ab}$, which is precisely what leads to (15).

As well as for the one-species case, this equation is valid only when ν_1 and ν_2 are small enough—in fact, both denominators have to be positive; one of them vanishing means that the critical filling is achieved for the corresponding species. In the special case (16), the condition is

$$\sqrt{\alpha_1}\nu_1 + \sqrt{\alpha_2}\nu_2 < \frac{1}{\sqrt{\max(\alpha_1, \alpha_2)}}. \quad (25)$$

4 Interpretation in terms of single-state distributions

We now wish to interpret the exactly solvable model of anyons in the LLL in terms of a gas of free particles with a particular statistical distribution. We will use arguments similar to those of [10] for the Calogero model. Description of single-state partition functions for multispecies systems follows Ref. [11].

Let there be single-particle states labeled with i , common for all species a , but with energies $\varepsilon_a^{(i)}$ possibly depending on a . The particles in different single-particle states are assumed to be statistically independent, which implies that the grand partition function is a product of single-state grand partition functions

$$\Xi = \prod_i \Xi^{(i)} \quad (26)$$

with $\Xi^{(i)}$ depending only on the Gibbs factors $x_a = e^{\beta(\mu_a - \varepsilon_a^{(i)})}$; $\Xi^{(i)} = \Xi^{(i)}(\{x_a\})$. The statistical distributions (average number of particles of different species in state i) are

$$n_a^{(i)} = x_a \frac{\partial}{\partial x_a} \ln \Xi^{(i)}. \quad (27)$$

We assume that $\ln \Xi^{(i)}$ may be expanded in integer powers of the Gibbs factors (again, we specify the formulas containing the series expansions for two species):

$$\ln \Xi^{(i)} = \sum_{k_1, k_2=0}^{\infty} f_{k_1 k_2} x_1^{k_1} x_2^{k_2}. \quad (28)$$

It follows then from (27)

$$n_a^{(i)} = \sum_{k_1, k_2=1}^{\infty} k_a f_{k_1 k_2} x_1^{k_1} x_2^{k_2}. \quad (29)$$

Combining (26) with (28), summing over single-particle states and comparing with (7), we get the relation

$$b_{k_1 k_2} = f_{k_1 k_2} \sigma_{k_1 k_2}, \quad (30)$$

where

$$\sigma_{k_1 k_2} = \sum_i e^{-\beta(k_1 \varepsilon_1^{(i)} + k_2 \varepsilon_2^{(i)})}. \quad (31)$$

We use the many-particle partition functions for anyons in the LLL in a harmonic well (19), with $\omega \rightarrow 0$, to calculate the coefficients $f_{k_1 k_2}$ in the expansion (28). Expressing the cluster coefficients in terms of the many-particle partition functions (but *not yet* using the thermodynamic limit prescription given in the Appendix), taking into account the expression (17) for the single-particle spectrum, in order to calculate $\sigma_{k_1 k_2}$, and then using (31) and (30), we eventually obtain

$$f_{k_1 0} = \frac{1}{k_1} \prod_{l_1=1}^{k_1-1} \left(1 - \frac{k_1 \alpha_1}{l_1}\right), \quad (32)$$

$$f_{k_1 k_2} = -\gamma \frac{k_1 + k_2}{k_1 k_2} \prod_{l_1=1}^{k_1-1} \left(1 - \frac{k_1 \alpha_1 + k_2 \gamma}{l_1}\right) \prod_{l_2=1}^{k_2-1} \left(1 - \frac{k_2 \alpha_2 + k_1 \gamma}{l_2}\right). \quad (33)$$

We thus conclude that in the limit $\omega \rightarrow 0$, multispecies anyons in the LLL can be described by the partition functions (28) and the statistical distributions (29) with the coefficients of their expansions in powers of the Gibbs factors given by (32)–(33).

Now we consider directly the case $\omega = 0$. All the single-particle states have then the same energy ω_c , therefore all the relevant quantities do not depend on the index i any longer, and summation over single-particle states reduces to multiplication by the number

of states in the LLL, $\rho_L V$, e. g. $\ln \Xi = \rho_L V \ln \Xi^{(1)}$. It turns out then that $\Xi^{(1)}$ can be represented as

$$\Xi^{(1)} = \prod_a \Xi_a \quad (34)$$

with

$$\Xi_a = 1 + \frac{n_a}{1 - \sum_b \alpha_{ab} n_b}, \quad (35)$$

so that for the coefficients of the expansions of the single-state grand partition functions

$$\Xi_a = \sum_{k_1, k_2} P_{a; k_1 k_2} x_1^{k_1} x_2^{k_2} \quad (36)$$

we obtain

$$P_{1; k_1 0} = \prod_{l_1=2}^{k_1} \left(1 - \frac{k_1 \alpha_1}{l_1} \right), \quad (37)$$

$$P_{1; k_1 k_2} = -\gamma \frac{k_1}{k_2} \prod_{l_1=2}^{k_1} \left(1 - \frac{k_1 \alpha_1 + k_2 \gamma}{l_1} \right) \prod_{l_2=1}^{k_2-1} \left(1 - \frac{k_2 \alpha_2 + k_1 \gamma}{l_2} \right) \quad (38)$$

and $P_{2; k_1 k_2} = P_{1; k_2 k_1} |_{\alpha_1 \leftrightarrow \alpha_2}$.

Eqs. (32) and (37) have been previously derived for fractional statistics in one dimension, starting from the Calogero model [10] (see also a recent derivation directly for exclusion statistics [15]). Eqs. (33) and (38) are generalizations onto the multispecies case.

The partition functions Ξ_a , in addition, satisfy the equations

$$\Xi_a = 1 + x_a \prod_b \Xi_b^{\delta_{ab} - \alpha_{ab}} \quad (39)$$

Substituting (35) into (39), we get the equation for the statistical distributions n_a

$$n_a / x_a = \prod_b (1 - \sum_c \alpha_{bc} n_c)^{\alpha_{ab}} (1 - n_a - \sum_c \alpha_{bc} n_c)^{\delta_{ab} - \alpha_{ab}}. \quad (40)$$

These statistical distributions coincide with the ones for exclusion statistics [8, 11] after the identification

$$g_{ab} = \alpha_{ab} \quad (41)$$

(in our model, consequently, the exclusion statistics parameters are symmetric, $g_{ab} = g_{ba}$).

It is now easy to calculate the cluster coefficients, using (30). Using the thermodynamic limit prescription at each order $k_1 + k_2$, (31) becomes

$$\sigma_{k_1 k_2} = \rho_L V e^{-\beta(k_1 + k_2)\omega_c} \quad (42)$$

and, with the use of (32)–(33), the formulas (21)–(22) are recovered. Note finally that once (34) and (35) have been correctly “guessed”, the equation of state is nothing but the thermodynamic identity $\beta PV = \ln \Xi$.

5 Concluding remarks

We have studied quantum and statistical mechanics for mutual fractional exchange statistics. We have presented a microscopic solvable model, anyons in the LLL, which, on one hand, involves mutual fractional exchange statistics and, on the other hand, reproduces the statistical mechanics for mutual fractional exclusion statistics. This establishes correspondence between the two approaches to mutual statistical interactions.

We hope that due to the similarity between anyons in the LLL and FQHE quasiparticles, the present work will help a better understanding of excitations in quantum Hall systems.

In addition, intimate links between anyons in the LLL and the Calogero model suggest that the microscopic solutions for multispecies anyons found in this paper can be used to construct solutions for a multispecies Calogero model which would reveal mutual exclusion statistics [16].

Acknowledgements: We thank Kåre Olaussen for valuable discussions. We are also grateful to NORDITA for their hospitality during the Anyon Workshop where this work

was initiated. S.M. gratefully acknowledges warm hospitality of the theory division of the IPN at Orsay where a significant part of this work was done and thanks Guillermo Zemba for bringing ref. [5] to his attention. S.B.I. was supported in part by the Russian Foundation for Fundamental Research under grant No. 95-02-04337.

Appendix

We summarize here, for reference purposes, the formulas concerning the cluster and virial expansions for a two-species gas, and give the relations between $Z_{N_1 N_2}$, $b_{n_1 n_2}$, and $a_{n_1 n_2}$, up to the fifth order.

The grand canonical partition function is

$$\Xi = \sum_{N_1, N_2} z_1^{N_1} z_2^{N_2} Z_{N_1 N_2} \quad (\text{A.1})$$

and the cluster coefficients $b_{k_1 k_2}$ are determined in terms of $Z_{N_1 N_2}$'s by writing the logarithm of (A.1) in the form

$$\ln \Xi = \sum_{k_1, k_2} b_{k_1 k_2} z_1^{k_1} z_2^{k_2}. \quad (\text{A.2})$$

The pressure and the densities are

$$\beta P = \frac{1}{V} \ln \Xi, \quad \rho_a = \frac{z_a}{V} \frac{\partial \ln \Xi}{\partial z_a}, \quad (\text{A.3})$$

respectively (V is the area); writing

$$\beta P = \sum_{k_1, k_2} a_{k_1 k_2} \rho_1^{k_1} \rho_2^{k_2}, \quad (\text{A.4})$$

using (A.3) with $\ln \Xi$ substituted from (A.2), and matching coefficients at equal powers of z 's on both sides, one gets the equations which allow to express $a_{k_1 k_2}$ in terms of $b_{k_1 k_2}$'s.

The explicit formulas are given below. For the cluster coefficients, one has

$$b_{10} = Z_{10};$$

$$b_{20} = -\frac{1}{2}Z_{10}^2 + Z_{20},$$

$$b_{11} = -Z_{01}Z_{10} + Z_{11};$$

$$b_{30} = \frac{1}{3}Z_{10}^3 - Z_{10}Z_{20} + Z_{30},$$

$$b_{21} = Z_{01}Z_{10}^2 - Z_{10}Z_{11} - Z_{01}Z_{20} + Z_{21};$$

$$b_{40} = -\frac{1}{4}Z_{10}^4 + Z_{10}^2Z_{20} - \frac{1}{2}Z_{20}^2 - Z_{10}Z_{30} + Z_{40},$$

$$b_{31} = -Z_{01}Z_{10}^3 + Z_{10}^2Z_{11} + 2Z_{01}Z_{10}Z_{20} - Z_{11}Z_{20} - Z_{10}Z_{21} - Z_{01}Z_{30} + Z_{31},$$

$$b_{22} = -\frac{3}{2}Z_{01}^2Z_{10}^2 + Z_{02}Z_{10}^2 + 2Z_{01}Z_{10}Z_{11} - \frac{1}{2}Z_{11}^2 - Z_{10}Z_{12} + Z_{01}^2Z_{20} \\ - Z_{02}Z_{20} - Z_{01}Z_{21} + Z_{22};$$

$$b_{50} = \frac{1}{5}Z_{10}^5 - Z_{10}^3Z_{20} + Z_{10}Z_{20}^2 + Z_{10}^2Z_{30} - Z_{20}Z_{30} - Z_{10}Z_{40} + Z_{50},$$

$$b_{41} = Z_{01}Z_{10}^4 - Z_{10}^3Z_{11} - 3Z_{01}Z_{10}^2Z_{20} + 2Z_{10}Z_{11}Z_{20} + Z_{01}Z_{20}^2 + Z_{10}^2Z_{21} \\ - Z_{20}Z_{21} + 2Z_{01}Z_{10}Z_{30} - Z_{11}Z_{30} - Z_{10}Z_{31} - Z_{01}Z_{40} + Z_{41},$$

$$b_{32} = 2Z_{01}^2Z_{10}^3 - Z_{02}Z_{10}^3 - 3Z_{01}Z_{10}^2Z_{11} + Z_{10}Z_{11}^2 + Z_{10}^2Z_{12} - 3Z_{01}^2Z_{10}Z_{20} \\ + 2Z_{02}Z_{10}Z_{20} + 2Z_{01}Z_{11}Z_{20} - Z_{12}Z_{20} + 2Z_{01}Z_{10}Z_{21} - Z_{11}Z_{21} \\ - Z_{10}Z_{22} + Z_{01}^2Z_{30} - Z_{02}Z_{30} - Z_{01}Z_{31} + Z_{32}.$$

If a harmonic oscillator regularization is used, the thermodynamic limit is meant as $\omega \rightarrow 0$, with a particular prescription at each order of the cluster expansion [17]. A treatment

analogous to the one given in [18] shows that to get the correct thermodynamic limit for $b_{k_1 k_2}$, the necessary prescription is to identify $\frac{2\pi}{(k_1+k_2)\beta m\omega^2} \rightarrow V$, in the limit $\omega \rightarrow 0$. For the virial coefficients, defining

$$\tilde{b}_{k_1 k_2} = b_{k_1 k_2} / b_{10}^{k_1} b_{01}^{k_2}, \quad (\text{A.5})$$

$$\tilde{a}_{k_1 k_2} = a_{k_1 k_2} / V^{k_1+k_2-1}, \quad (\text{A.6})$$

one has

$$\tilde{a}_{10} = 1;$$

$$\tilde{a}_{20} = -\tilde{b}_{20},$$

$$\tilde{a}_{11} = -\tilde{b}_{11};$$

$$\tilde{a}_{30} = 4\tilde{b}_{20}^2 - 2\tilde{b}_{30},$$

$$\tilde{a}_{21} = \tilde{b}_{11}^2 + 4\tilde{b}_{11}\tilde{b}_{20} - 2\tilde{b}_{21};$$

$$\tilde{a}_{40} = -20\tilde{b}_{20}^3 + 18\tilde{b}_{20}\tilde{b}_{30} - 3\tilde{b}_{40},$$

$$\tilde{a}_{31} = -\tilde{b}_{11}^3 - 6\tilde{b}_{11}^2\tilde{b}_{20} - 24\tilde{b}_{11}\tilde{b}_{20}^2 + 3\tilde{b}_{11}\tilde{b}_{21} + 12\tilde{b}_{20}\tilde{b}_{21} + 9\tilde{b}_{11}\tilde{b}_{30} - 3\tilde{b}_{31},$$

$$\begin{aligned} \tilde{a}_{22} = & -9\tilde{b}_{02}\tilde{b}_{11}^2 - 3\tilde{b}_{11}^3 + 6\tilde{b}_{11}\tilde{b}_{12} - 12\tilde{b}_{02}\tilde{b}_{11}\tilde{b}_{20} - 9\tilde{b}_{11}^2\tilde{b}_{20} + 6\tilde{b}_{12}\tilde{b}_{20} + 6\tilde{b}_{02}\tilde{b}_{21} \\ & + 6\tilde{b}_{11}\tilde{b}_{21} - 3\tilde{b}_{22}; \end{aligned}$$

$$\tilde{a}_{50} = 112\tilde{b}_{20}^4 + 32\tilde{b}_{40}\tilde{b}_{20} + 18\tilde{b}_{30}^2 - 144\tilde{b}_{30}\tilde{b}_{20}^2 - 4\tilde{b}_{50},$$

$$\begin{aligned} \tilde{a}_{41} = & \tilde{b}_{11}^4 + 8\tilde{b}_{11}^3\tilde{b}_{20} + 40\tilde{b}_{11}^2\tilde{b}_{20}^2 + 160\tilde{b}_{11}\tilde{b}_{20}^3 - 4\tilde{b}_{11}^2\tilde{b}_{21} - 24\tilde{b}_{11}\tilde{b}_{20}\tilde{b}_{21} - 80\tilde{b}_{20}^2\tilde{b}_{21} \\ & + 2\tilde{b}_{21}^2 - 12\tilde{b}_{11}^2\tilde{b}_{30} - 120\tilde{b}_{11}\tilde{b}_{20}\tilde{b}_{30} + 24\tilde{b}_{21}\tilde{b}_{30} + 4\tilde{b}_{11}\tilde{b}_{31} + 24\tilde{b}_{20}\tilde{b}_{31} \end{aligned}$$

$$\begin{aligned}
& + 16\tilde{b}_{11}\tilde{b}_{40} - 4\tilde{b}_{41}, \\
\tilde{a}_{32} = & 16\tilde{b}_{02}\tilde{b}_{11}^3 + 6\tilde{b}_{11}^4 - 12\tilde{b}_{11}^2\tilde{b}_{12} + 48\tilde{b}_{02}\tilde{b}_{11}^2\tilde{b}_{20} + 32\tilde{b}_{11}^3\tilde{b}_{20} - 32\tilde{b}_{11}\tilde{b}_{12}\tilde{b}_{20} \\
& + 64\tilde{b}_{02}\tilde{b}_{11}\tilde{b}_{20}^2 + 80\tilde{b}_{11}^2\tilde{b}_{20}^2 - 32\tilde{b}_{12}\tilde{b}_{20}^2 - 24\tilde{b}_{02}\tilde{b}_{11}\tilde{b}_{21} - 20\tilde{b}_{11}^2\tilde{b}_{21} + 8\tilde{b}_{12}\tilde{b}_{21} \\
& - 32\tilde{b}_{02}\tilde{b}_{20}\tilde{b}_{21} - 64\tilde{b}_{11}\tilde{b}_{20}\tilde{b}_{21} + 8\tilde{b}_{21}^2 + 8\tilde{b}_{11}\tilde{b}_{22} + 16\tilde{b}_{20}\tilde{b}_{22} - 24\tilde{b}_{02}\tilde{b}_{11}\tilde{b}_{30} \\
& - 24\tilde{b}_{11}^2\tilde{b}_{30} + 12\tilde{b}_{12}\tilde{b}_{30} + 8\tilde{b}_{02}\tilde{b}_{31} + 12\tilde{b}_{11}\tilde{b}_{31} - 4\tilde{b}_{32};
\end{aligned}$$

with the obvious symmetry with respect to permutation of the subscripts.

References

- [1] J.M. Leinaas and J. Myrheim, Nuovo Cim. **37B** (1977) 1.
- [2] F. Wilczek, Phys. Rev. Lett. **48** (1982) 1144; *ibid.* **49** (1982) 957.
- [3] P.W. Anderson, Science **235** (1987) 1196; G. Baskaran et al., Solid State Commun. **63** (1987) 973; T. Mor, Phys. Lett. A **184** (1993) 29.
- [4] F. Wilczek, Phys. Rev. Lett. **69** (1992) 132; Z. F. Ezawa and A. Iwazaki, Phys. Rev. B **47** (1993) 7295 and references therein.
- [5] J. Fröhlich, A. Zee, Nucl. Phys. **B364** (1991) 517.
- [6] A. Dasnières de Veigy and S. Ouvry, Phys. Rev. Lett. **72** (1994) 600.
- [7] A. Dasnières de Veigy and S. Ouvry, Mod. Phys. Lett. B **9** (1995) 271.
- [8] Y.-S. Wu, Phys. Rev. Lett. **73** (1994) 922.
- [9] F.D.M. Haldane, Phys. Rev. Lett. **67** (1991) 937.
- [10] S.B. Isakov, Int. J. Mod. Phys. A **9** (1994) 2563.

- [11] S.B. Isakov, Mod. Phys. Lett. B **8** (1994) 319.
- [12] L. Brekke, A.F. Falk, S.J. Hugen and T.D. Imbo, Phys. Lett. A **271** (1991) 73.
- [13] A. Dasnières de Veigy and S. Ouvry, Phys. Lett. **B307** (1992) 91.
- [14] D.P. Arovas, J.R. Schrieffer, F. Wilczek, A. Zee, Nucl. Phys. **B251** (1985) 117.
- [15] A. Polychronakos, preprint UUITP-03/95, hep-th/9503077.
- [16] C. Furtlehner and S. Ouvry, IPNO/TH 94-02, to be published in MPLB
- [17] A. Comtet, Y. Georgelin and S. Ouvry, J. Phys. A (Math.Gen.) **22** (1989) 3917;
J. McCabe and S. Ouvry, Phys. Lett. **B260** (1991) 113.
- [18] K. Olaussen, Trondheim preprint No. 13 (1992).